arXiv:1303.0764v1 [cond-mat.str-el] 4 Mar 2013

Electric polarization in correlated insulators

R. Nourafkan, and G. Kotliar
Department of Physics & Astronomy, Rutgers University, Piscataway, NJ 08854-8019, USA

We derive a formula for the electric polarization of interacting insulators, expressed in terms of the full Green's and vertex functions of the system. We use the formula to investigate changes in the electric polarization of the half-filled ionic Hubbard model in the DMFT framework. Interaction renormalizes both ionicity, and covalency of the system and competition between these renormalized quantities determines the electric polarization of a correlated band insulator. Based on a quasi-particle approximation, we comment that the LDA+U based calculations, which *does not account* for the quasi-particle weight renormalization, will overestimate the electronic part of the electric polarization. This is sizable when the quasi-particle weight is far from one for active bands in the polarization.

PACS numbers: 77.84.-s, 77.22.Ej, 03.65.Vf

INTRODUCTION

The electric polarization, $\mathbf{P}$, is a measure of the position differences between the center of the band electrons and the lattice ions. It is generally non zero for a material which lacks inversion symmetry and plays the role of the order parameter in the theory of ferroelectricity. Over the past two decades, there have been important conceptual and computational advances in the first principles calculation of this quantity. [1]

The modern theory of polarization expresses $\mathbf{P}$ in terms of the Berry phase acquired by a Slater determinant, describing the insulating state as an effective one particle theory. [2] A complementary picture of the polarization, as the displacement of the Wannier centers of the single particle states involve, emerges naturally in this formalism. [3]

The Berry phase formalism was extended beyond an effective single particle picture, by considering the changes of the phase of the many body function of all the electrons in the solid as a response to the changes in the boundary conditions. [4, 5] The resulting expression can also be recast in terms of the resolvent of the Hamiltonian of all the interacting electrons in the many body ground state. [6] The many body wave function formalism to calculate polarization was applied to simple model Hamiltonians, [7] but is prohibitively difficult to carry out in practice for realistic models of the electronic structure of a solid.

In this paper we formulate the problem of the calculation of the electronic polarization of a correlated electronic system in terms of the one particle Greens function and vertex functions. We derive a general expression for the polarization, and discuss its implementation within dynamical mean field theory (DMFT). [8] We apply the formalism to a simple model of a correlated insulator. The approach has similarities to recent studies of incorporating correlations in the calculation of the topological indices of topological insulators [9, 10], and to the early work of Volovik [11] and gives useful insights into the effects of Hubbard correlations on the electric polarization.

In an extended system the change in polarization, $\Delta\mathbf{P}$, induced by distorting an atomic coordinate, $\mathbf{R}_{i\alpha} \to \mathbf{R}_{i\alpha} + \xi\Delta\mathbf{R}_{i\alpha}$, that breaks inversion symmetry is given by the integrated bulk transient current as the system adiabatically evolves from the initial state ($\xi = 0$) to the final state ($\xi = 1$), i.e., [1]

$$\begin{aligned}\Delta\mathbf{P} &= \int_0^1 d\xi\, \frac{d\mathbf{P}}{d\xi} = \int_0^1 d\xi\, \frac{1}{V_{\rm cell}} \int_{\rm cell} d\mathbf{r}\, \mathbf{J}(\mathbf{r},\xi) \\ &= \int_0^1 d\xi\, \mathbf{J}(\mathbf{q}=\mathbf{0},\xi) \qquad (1)\end{aligned}$$

where $\mathbf{J}(\mathbf{r},\xi)$ denotes the current density and $V_{\rm cell}$ is the primitive-cell volume. Implicit in the analysis is that the system must remain insulating everywhere along the path, as otherwise the adiabatic condition fails.

Assume the system is described by $\hat{H}(\xi)$ for a given value of the distortion. A small change in ξ, i.e., $\xi \to \xi + \delta\xi$, perturbs the Hamiltonian of the system by $(\delta\hat{H}/\delta\xi)\delta\xi \equiv \hat{H}'_\xi\delta\xi$ and induces a current. In order to evaluate the current explicitly, we assume $\delta\xi(t) = \delta\xi\exp(-i\nu t + \eta t)$ with η an infinitesimal positive number and finally we consider the $\nu \to 0$ limit.

According to the Kubo formula the current density induced by a perturbation is $\mathbf{J}(\nu) = C^{(\xi)R}_{\mathbf{j}H'}(\mathbf{q} = \mathbf{0},\nu)\delta\xi$, where $C^{(\xi)R}_{\mathbf{j}H'}(\mathbf{q} = \mathbf{0},\nu)$ is $\mathbf{q} = \mathbf{0}$ component of the Fourier transform (in space and time) of the retarded correlation function, [12]

$$C^{(\xi)R}_{\mathbf{j}(\mathbf{r})H'(\mathbf{r}')}(t-t') = -i\Theta(t-t')\langle[\hat{\mathbf{j}}_\xi(\mathbf{r},t),\hat{H}'_\xi(\mathbf{r}',t')]\rangle_\xi. \qquad (2)$$

Here the subscript ξ for operators shows that they are in the Heisenberg representation with respect to the instantaneous Hamiltonian and $\langle\cdots\rangle_\xi$ means that the average is taken with respect to the instantaneous spectrum. In the limit of $\nu = 0$ (static distortion), $C^{(\xi)R}_{\mathbf{j}H'}(\mathbf{q} = \mathbf{0},\nu = 0) = 0$, therefore the driving force is $\delta\dot{\xi}$ and $\mathbf{J} \propto \delta\dot{\xi}$. We evaluate $\mathbf{J}$ as

$$\begin{aligned}\mathbf{J}(\mathbf{q}=\mathbf{0},\xi) &= i\Big(\frac{\partial}{\partial\nu}C^{(\xi)R}_{\mathbf{j}H'}(\mathbf{q}=\mathbf{0},\nu)\Big)\Big|_{\nu=0}\delta\dot{\xi} \\ &= i\Big(\frac{\partial}{\partial i\nu_n}C^{(\xi)}_{\mathbf{j}H'}(\mathbf{q}=\mathbf{0},i\nu_n)\Big)\Big|_{\nu_n=0}\delta\dot{\xi} \equiv \mathcal{L}^{(\xi)}_{\mathbf{j}H'}\delta\dot{\xi} \qquad (3)\end{aligned}$$

where we used analytic continuation in the last line and we defined transport coefficient $\mathcal{L}^{(\xi)}_{\mathbf{j}H'}$. Substituting Eq. (3) in Eq. (1) gives an expression for $\Delta\mathbf{P}$.

Note that $\Re\mathcal{L}^{(\xi)}_{\mathbf{j}H'}$, is anti-symmetric under exchange of $\mathbf{j}$ and H' operators, i.e., $\Re\mathcal{L}^{(\xi)}_{\mathbf{j}H'} = -\Re\mathcal{L}^{(\xi)}_{H'\mathbf{j}}$, which can be seen from its Lehmann representation (see supplemental material). [13]

In a diagrammatic series expansion we can express $C^{(\xi)R}_{\mathbf{j}H'}$ in terms of interacting Green's functions and irreducible particle-hole vertex functions. We can find the perturbation expansion for $\mathcal{L}_{\mathbf{j}H'}$ by differentiating with respect to $i\nu_n$. This introduces a derivative vertex $(\partial/\partial i\omega_m)$ (see supplemental material). [13] The diagrams contributing at the $\mathcal{L}_{\mathbf{j}H'}$ are of two types: (i) diagrams which are separated into three pieces, each of which has a vertex, by cutting three electron lines. We denote the contribution of these diagrams to the polarization by $\Delta\mathbf{P}_1$. (ii) All the rest diagrams which their contribution to the polarization is denoted by $\Delta\mathbf{P}_2$. The change in polarization from the diagrams of type (i) is given by

$$\Delta\mathbf{P}_1 = -\int_0^1 d\xi\,\left(\frac{ie}{2N\beta}\right)\sum_{\mathbf{k}\sigma}\sum_{\omega_m} \mathrm{Tr}\left(\mathbf{\Lambda}^{(\xi)}_{\mathbf{j}}\mathbf{G}^{(\xi)}\mathbf{\Lambda}^{(\xi)}_{H'}\mathbf{G}^{(\xi)}\frac{\partial\mathbf{G}^{(\xi)-1}}{\partial i\omega_m}\mathbf{G}^{(\xi)} - \mathbf{\Lambda}^{(\xi)}_{\mathbf{j}} \leftrightarrow \mathbf{\Lambda}^{(\xi)}_{H'}\right) \quad (4)$$

where we have used the identity $(\partial\mathbf{G}^{(\xi)}/\partial i\omega_m) = -\mathbf{G}^{(\xi)}(\partial\mathbf{G}^{(\xi)-1}/\partial i\omega_m)\mathbf{G}^{(\xi)}$ and the anti-symmetric property of $\mathcal{L}_{\mathbf{j}H'}$ in writing Eq. (4) down. $\mathbf{\Lambda}_{\mathbf{j}}$, $\mathbf{\Lambda}_{H'}$ are the particle-hole vertex functions and β is the inverse temperature. The interacting single particle Green's function of the system is

$$\mathbf{G}^{(\xi)}_{\mathbf{k}\sigma}(i\omega_m) = [(i\omega_n+\mu)\mathbf{1} - \mathbf{H}^{(\xi)}_0(\mathbf{k}) - \mathbf{\Sigma}^{(\xi)}_{\sigma}(\mathbf{k}, i\omega_m)]^{-1}, \quad (5)$$

where $\mathbf{H}^{(\xi)}_0(\mathbf{k})$ denotes the non-interacting part of the Hamiltonian. While $\mathbf{G}^{(\xi)}$ and $\mathbf{H}^{(\xi)}_0$ are gauge covariant, $\Delta\mathbf{P}$ is gauge invariant. However, the select of the interaction matrix and the expression of the current do depend on the gauge. [14] Here we write $\mathbf{H}^{(\xi)}_0$ in the *periodic gauge*. [15] $\mathbf{\Sigma}^{(\xi)}_{\sigma}(\mathbf{k}, i\omega_m)$ denotes the electron self-energy. Bold quantities are $n\times n$ matrices where n denotes number of orbitals within the unit cell.

One can use the Ward identity to replace the electric current vertex by $\mathbf{\Lambda}^{(\xi)}_{\mathbf{j}}(\mathbf{k}) = -(\partial\mathbf{G}^{(\xi)}_{\mathbf{k}\sigma}(i\omega_m)^{-1}/\partial\mathbf{k})$ and the distortion current vertex by $\mathbf{\Lambda}^{(\xi)}_{H'}(\mathbf{k}) = -(\partial\mathbf{G}^{(\xi)}_{\mathbf{k}\sigma}(i\omega_m)^{-1}/\partial\xi)$. [16] Therefore, the Eq. (4) can be rewritten as

$$\Delta\mathbf{P}_1 = -\int_0^1 d\xi\,\left(\frac{ie}{N\beta}\right)\sum_{\mathbf{k}\sigma}\sum_{\omega_m}\frac{1}{6}\varepsilon^{\mathbf{k}\xi\omega_m} \mathrm{Tr}\left(\frac{\partial\mathbf{G}^{(\xi)-1}}{\partial\mathbf{k}}\mathbf{G}^{(\xi)}\frac{\partial\mathbf{G}^{(\xi)-1}}{\partial\xi}\mathbf{G}^{(\xi)}\frac{\partial\mathbf{G}^{(\xi)-1}}{\partial i\omega_m}\mathbf{G}^{(\xi)}\right) \quad (6)$$

where we have introduced an anti-symmetric tensor, $\varepsilon^{\mathbf{k}\xi\omega_m}$. Eq. (6) follows from Eq. (4) due to the anti-symmetric property of the derivative relative to $\mathbf{k}$ and ξ and cyclic property of the trace. In the non-interacting case Eq. (6) reduces to the Berry phase formula (see supplemental material). [2, 13]

It is not possible to write $\Delta\mathbf{P}_2$ in the form of Eq. (4). However, we have evaluated it perturbatively in various models and found it to be very small.

In the following we will work in the DMFT approximation, where we ignore the current vertex corrections while keeping the self-energy corrections associated with electron correlation. In DMFT, $\Delta\mathbf{P}_2$ vanishes and the correlation function reduces to the bubble contribution with the fully interacting Green's function and the following vertices; $(\partial\mathbf{G}^{(\xi)-1}/\partial\mathbf{k}) = -(\partial\mathbf{H}^{(\xi)}_0/\partial\mathbf{k})$, $(\partial\mathbf{G}^{(\xi)-1}/\partial\xi) = -(\partial\mathbf{H}^{(\xi)}_0/\partial\xi) - (\partial\mathbf{\Sigma}^{(\xi)}/\partial\xi)$, and $(\partial\mathbf{G}^{(\xi)-1}/\partial i\omega_n) = \mathbf{1} - (\partial\mathbf{\Sigma}^{(\xi)}/\partial i\omega_n)$.

To investigate the influence of Hubbard correlations on the electronic part of the polarization and to benchmark our formalism we turn to a simple model of ferroelectric materials, the ionic Hubbard model (IHM) in one dimension. On a bipartite lattice the IHM Hamiltonian is defined by the following equation [17–20]:

$$H = -t\sum_{i\sigma}[1+(-1)^i\xi](c^{\dagger}_{i\sigma}c_{i+1\sigma} + H.c.) + \Delta\sum_{i\sigma}(-1)^i n_{i\sigma} + U\sum_i n_{i\uparrow}n_{i\downarrow} \quad (7)$$

where, $t(>0)$ and U denote the hopping amplitude between nearest-neighbor sites and the on-site Coulomb repulsion, respectively. $-\Delta(+\Delta)$ denotes the local potential energy for the $A(B)$ sublattice, respectively.

The model is relevant for the study of organic ferroelectrics with stacks of alternating donor and acceptor molecules. The paraelectric phase is centrosymmetric (so that all $A-B$ distances are the same), whereas in the ferroelectric phase there is a relative displacement of the A and B sublattices, which breaks the inversion symmetry and produces an alternating pattern of short and long $A-B$ distances. Here we modify the hopping amplitude by $\pm\xi$ alternatively to include a dimer term that breaks the inversion symmetry.

At half filling, the noninteracting undistorted system ($\xi = 0$) is a band insulator (BI) with a charge gap equal to $\delta_c = 2\Delta$. Upon turning on and increasing U the charge gap, δ_c, in the correlated band insulator shrinks until a discontinuous transition to a Mott insulator (MI) occurs with a hysteresis region, if the anti-ferromagnetic (AF) long-range order is not allowed to set in. In the Mott phase a large gap is established. [21, 22] As the system is driven deeper into the Mott phase, the gap size increases. Both the BI and the MI are non-polar at $\xi = 0$; however, depending on how one defines the zero reference point of the polarization one can assign a polarization value to these states. Here we define the zero reference point to be the usual band insulator where both electrons occupy the Wannier function centered on the lower energy site, then using the classical point charge model we assign $P^{(\mathrm{el})}/ea = -1/2$ to a Mott insulator with no long range order. In the centrosymmetric system without spontaneous inversion symmetry breaking [24], increasing U from zero causes a charge exchange between sites. However, charge flows symmetrically, the macro-



FIG. 1. (Color online) Electric polarization as a function of distortion, ξ, for several value of U in the PM phase (For $U/t < 7.5$ the system is in the band insulator phase with zero reference point of polarization $P(\xi = 0)/ea = 0$. For $U/t = 8.0$ data is shown in the Mott phase with zero reference point of polarization $P(\xi = 0)/ea = -1/2$). Black dashed lines show Eq. (13).

scopic current is zero and therefore $\Delta P^{(\mathrm{el})} = 0$. In the presence of the AF order, system has a transition to the AF phase at a smaller critical U. [23]

Next we present our results in the presence of a distortion. We set $\Delta = 0.5t$ and use Eq. (6) and DMFT to evaluate the electronic part of the electric polarization of the distorted IHM in the paramagnetic (PM) and AF phases. The Bloch representation of the non-interacting part of the IHM Hamiltonian, Eq. (7), is given by

$$\mathbf{H}_0(k) = -2t\cos(ka/2)\boldsymbol{\tau}_x + 2\xi t\sin(ka/2)\boldsymbol{\tau}_y - \Delta\boldsymbol{\tau}_z \quad (8)$$

where $\boldsymbol{\tau}_{x,y,z}$ are the Pauli matrices and a is unit cell length. In the DMFT approximation the current vertex and the distortion vertex are given by

$$\frac{\partial \mathbf{G}^{(\xi)-1}}{\partial k} = at\sin(ka/2)\boldsymbol{\tau}_x - a\xi t\cos(ka/2)\boldsymbol{\tau}_y, \quad (9)$$

$$\frac{\partial \mathbf{G}^{(\xi)-1}}{\partial \xi} = -2t\sin(ka/2)\boldsymbol{\tau}_y - \frac{\partial \boldsymbol{\Sigma}^{(\xi)}}{\partial \xi}. \quad (10)$$

where $\boldsymbol{\Sigma}$ is a diagonal matrix with elements Σ_A and Σ_B.

Fig. 1 shows the polarization of the system, $P^{(\mathrm{el})}_{\Delta,U}(\xi)/ea$, obtained from an evaluation of Eq. (6) for several values of U as a function of ξ. As can be seen in the correlated BI ($U < U_c \simeq 7.5t$), the absolute value of the $P^{(\mathrm{el})}_{\Delta,U}(\xi)/ea$ increases with ξ and saturates at $-1/2$. The saturation value happens at smaller ξ upon increasing U (note that for small values of U the saturation value is not apparent on this figure, as it occurs at larger ξ). In the non-interacting case, a system with small ionicity is more polarizable. In order to explain results with non-zero U with $U < U_c$, one needs to understand how the correlation renormalizes the ionicity and the covalency of the system. To get some insight we work in the QP approximation and derive an analytical equation for the electric polarization at small dimerization. The QP Green's function is given by

$$\mathbf{G}^{(\xi)}_{qp}(\mathbf{k},\omega) = \mathbf{z}^{1/2}[i\omega_n\mathbf{1} - \mathbf{H}^{(\xi)}_{qp}(\mathbf{k})]^{-1}\mathbf{z}^{1/2}, \quad (11)$$

where $\mathbf{H}^{(\xi)}_{qp}(\mathbf{k}) = \mathbf{z}^{1/2}(\mathbf{H}^{(\xi)}_0(\mathbf{k}) + \Re\boldsymbol{\Sigma}^{(\xi)}_U(0) - \mu\mathbf{1})\mathbf{z}^{1/2}$ is the QP Hamiltonian. The quasi-particle weight is denoted by $\mathbf{z} \simeq [\mathbf{1} - \Im\boldsymbol{\Sigma}(i\omega_0)/\omega_0]^{-1}$, where $\omega_0 = \pi/\beta$ is first fermionic Matsubara frequency. Substituting Eq. (11) into Eq. (6) and using the fact that $\partial\mathbf{G}^{-1}_{qp}/\partial i\omega_n = \mathbf{z}^{-1}$, one can see that an extension of Eq. (S13) in the supplemental material [13] to the interacting IHM is obtained by replacing bare quantities with renormalized ones, $\Delta \to \Delta^{ren}_{qp}$ and $t \to zt$. In our case $\mathbf{z}$ is a diagonal matrix with $z_A = z_B \equiv z$. The renormalized ionicity can be obtained from the renormalized charge gap which in turn can be obtained by identifying the poles of the renormalized propagator, Eq. (11), and is given by

$$\delta^{qp}_c = 2\Delta^{ren}_{qp} = 2z[\Delta - \Re(\Sigma_A(i\omega_0) - \Sigma_B(i\omega_0))]. \quad (12)$$

The frequency range of the validity of the Fermi liquid assumptions decreases upon increasing U, but the charge gap also shrinks simultaneously. A closer investigation of the self-energy and local density states of the correlated insulator show that one can continue to assume the Fermi liquid assumptions on the frequency range between the highest pole of the valance band and the lowest pole of the conduction band for all interaction strengths. Therefore, Eq. (11) describes low energy physics precisely and the Eq. (12) is the true charge excitation gap, $\delta_c = \delta^{qp}_c$.

Thus, in the QP approximation the polarization for small distortions is given by:

$$P^{(\mathrm{el})}_{\Delta,U}(\xi) \simeq -\frac{2ea}{N}\sum_k \frac{(\Delta^{ren}_{qp}/(zt))\sin^2(ka/2)}{[(\Delta^{ren}_{qp}/(zt))^2 + 4\cos^2(ka/2)]^{3/2}}\xi \quad (13)$$

which shows that for small ξ the electric polarization of the interacting system is the same as a non-interacting system with effective ionicity given by the charge gap of the system renormalized by quasi-particle weight, $(\Delta^{ren}_{qp}/(zt))$. Upon increasing U, (Δ^{ren}_{qp}/t) decreases faster than z which leads to a reduction of $(\Delta^{ren}_{qp}/(zt))$ which in turn leads to a more polarizable system where a small distortion can trigger substantial changes in the electric polarization. In Fig. 1 the QP result Eq. (13) is shown with dashed lines. The agreement between the QP approximation and the full answer is very good in the linear regime of polarization. Notice that at strong interaction strengths Eq. (13) fails because it is built on the assumptions that $\Re\boldsymbol{\Sigma}$ is linear and $\Im\boldsymbol{\Sigma}$ vanishes identically on the whole bandwidth.

Fig. 2 shows $P^{(\mathrm{el})}_{\Delta,U}(\xi = 0.08t)/ea$ for several values of U obtained from the full- and QP-calculations. In the QP-calculation we employ $\mathbf{H}^{(\xi)}_{qp}$ to define an effective non-interacting system and calculate P using the Berry phase formula. [2] It is worth mentioning that same results for polarization can be obtained by working with $(\mathbf{H}^{(\xi)}_0(\mathbf{k}) + \Re\boldsymbol{\Sigma}^{(\xi)}_U(0) - \mu\mathbf{1})$ as the QP Hamiltonian, which provides the correct ratio of the renormalized ionicity to the covalency. However, the latter Hamiltonian gives incorrect results on the charge gap and does not account for quasi-particle weight renormalization. Consistently with Eq. (13), the QP results agree very



FIG. 2. (Color online) Electric polarization for a distorted system with $\xi = 0.08t$ for several interaction strengths, U, using the full calculation and the quasi-particle (QP) approximation.

well with the full calculations at weak to intermediate interaction strengths. However, at strong coupling, P obtained from the QP method deviates from the full answer because it does not account for the bandwidth renormalization correctly. Inset of the Fig. 2 shows the charge gap as a function of U.

At large interaction strength, for example $U = 8.0t$, the system is in a Mott phase. In a Mott Phase, the variation in the electronic part of the polarization as a function of the distortion is very small (here we restrict ourself to small ξ). Indeed in a Mott phase, charge fluctuations are strongly suppressed and an ionic displacement induces only a very small current. In the Mott phase the effective non-interacting system described by $\mathbf{H}_{qp}$ is an extremely ionic system where one sublattice is fully occupied and the other one is empty. In reality, however, in the true ground state the charge is almost uniformly distributed in the system. Thus, the QP approximation fails in the Mott phase.

Next we investigate the AF phase which allows us to compare our results with the QMC results on IHM. Investigation of the staggered magnetization, $m = \langle n_{A\uparrow} - n_{A\downarrow}\rangle = -\langle n_{B\uparrow} - n_{B\downarrow}\rangle$, of the centrosymmetric structure, $\xi = 0$, as a function of the interaction strength shows that at zero temperature the system shows long-range AF for $U \geq 2.4t$. The onset of the AF phase coincides with the onset of the possible spontaneous bond order phase. [25] Fig. 3 shows $|P^{(\mathrm{el})}_{\Delta,U}(\xi = 0.02) - P^{(\mathrm{el})}_{\Delta,U}(\xi = 0)|$ obtained from an implementation of Eq. (6) in the AF phase along with a QMC calculation reproduced from [7]. In the QMC data, the phase diagram of the undistorted ionic Hubbard model includes a spontaneous dimerized phase with non-zero polarization which means that the reference point of polarization, $P^{(\mathrm{el})}_{\Delta,U}(\xi = 0)$, has different value than us. However, the shown data is ΔP which is independent of the reference point. Also, the QMC calculation was done for a slightly different Δ. Nevertheless, the two calculations agree very well with one another.

In conclusion, we have introduced a practical many body approach to the calculation of the electric polarization of interacting insulators. We have implemented and tested the formalism in the context of a model Hamiltonian and shown that while the electronic polarization is affected by Hubbard correlations, it is a less sensitive quantity well approximated by a quasi-particle approximation and depending on the ratio of the gap and the covalency of the quasi-particle band structure. These results justifying the success of the traditional electronic structure of methods which do not include an explicit frequency dependence of the self energy . Notice however, that in some correlated materials such as HoMnO$_3$ this method has been reported to overestimate the electric polarization. [26, 27] This can be understood as resulting from the effects of the bandwidth renormalization on the polarization described here. The formalism of this paper can be implemented in realistic LDA+DMFT codes. This together with a comparison with accurate experimental ARPES studies of correlated ferroelectrics are interesting open problems.



FIG. 3. (Color online) $|P^{(\mathrm{el})}_{\Delta,U}(\xi = 0.02) - P^{(\mathrm{el})}_{\Delta,U}(\xi = 0)|$ obtained from an implementation of Eq. (6) in the AF phase along with a QMC calculation reproduced from [7]. Inset shows a schematic phase diagram of the system.

We are grateful to F. Marsiglio, M. E. Pezzoli, A. Soluyanov, G. Ortiz, K. Haule and D. Vanderbilt for useful discussions. This work has been supported by NSF DMR-0906943.

[1] R. Resta and D. Vanderbilt, *Physics of Ferroelectrics: a Modern Perspective*, Springer-Verlag (2007).
[2] R. D. King-Smith and D. Vanderbilt, Phys. Rev. B **47**, 1651 (1993).
[3] D. Vanderbilt and R. D. King-Smith, Phys. Rev. B **48**, 4442 (1993).
[4] G. Ortiz, and R. M. Martin, Phys. Rev. B **49**, 14202 (1994).
[5] R. Resta, Phys. Rev. Lett. **80**, 1800 (1998).
[6] G. Oritz, private communication.
[7] T. Wilkens, and R. M. Martin, Phys. Rev. B **63**, 235108 (2001).
[8] A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg Rev. Mod. Phys. **68**, 13 (1996).
[9] Z. Wang, X. L. Qi, and S. C. Zhang, Phys. Rev. Lett. **105**, 256803 (2010).
[10] A. Go, W. W. Krempa, G. S. Jeon, K. Park, and Y. B. Kim, Phys. Rev. Lett. **109**, 066401 (2012).
[11] G. E. Volovik, *The Universe in a Helium Droplet* (Oxford, 2003).
[12] H. Bruus and K. Flensberg, *Many-Body Quantum Theory in*

Condensed Matter Physics: An Introduction , Oxford University Press (2004).

[13] See Supplemental Material at http://... for a discussion of the diagrammatic expansion of the correlation function and reduction of the Eq. (4) to Berry phase formula at the non-interacting limit.

[14] I. Paul and G. Kotliar, Phys. Rev. B **67**, 115131 (2003).

[15] In the periodic gauge, $\mathbf{H}_0$ satisfies $\mathbf{H}_0(\mathbf{k}+\mathbf{K}) = O^\dagger\mathbf{H}_0(\mathbf{k})O$, where O is a diagonal matrix with $O_{nn} = \exp(i\mathbf{K}.\mathbf{d}_n)$, n-th orbital is located at $\mathbf{d}_n$, and $\mathbf{K}$ is the reciprocal lattice vector [K. T. Chen, P. A. Lee, Phys. Rev. B **84**, 205137 (2011)].

[16] K. Ishikawa and T. Matsuyama, Nucl. Phys. B **280**, 523 (1987).

[17] G. Giovannetti, S. Kumar, A. Stroppa, J. V. D. Brink, and S. Picozzi, Phys. Rev. lett., **103**, 266401 (2009).

[18] T. Egami, S. Ishihara and M. Tachiki, Science **261**, 1307 (1993); Phys. Rev. B **49**, 8944 (1994).

[19] R. Resta and S. Sorella, Phys. Rev. Lett. **74**, 4738 (1995).

[20] L. D. Freo, A. Painelli, and Z. G. Soos, Phys. Rev. Lett., **89**, 027402 (2002).

[21] A. Garg, H.R. Krishnamurthy, and M. Randeria, Phys. Rev. Lett. **97**, 046403 (2006).

[22] L. Craco, P. Lombardo, R. Hayn, G. Japaridze, E. Hartmann, Phys. Rev. B **78**, 075121 (2008).

[23] K. Byczuk, M. Sekania, W. Hofstetter, A. Kampf, Phys. Rev. B **79**, 121103(R) (2009).

[24] In a centrosymmetric system of interacting electrons, the variation of the polarization as a function of U depends on whether spontaneous breaking of inversion symmetry happens or not. A DMFT calculation cannot capture a spontaneous inversion symmetry broken phase and Cluster-DMFT breaks translational symmetry by construction, so for the centrosymmetric system we only consider the symmetric Mott phase.

[25] S. S. Kancharla and E. Dagotto, Phys. Rev. Lett., **98**, 016402 (2007).

[26] S. Picozzi, K. Yamauchi, B. Sanyal, I. A. Sergienko, and E. Dagotto, Phys. Rev. Lett. **99**, 227201 (2007).

[27] Y. S. Chai, *et all*, Phys. Rev. B **85**, 184406 (2012).

Electric polarization in a correlated insulator - Supplemental materials

I. ANTI-SYMMETRIC PROPERTY

The anti-symmetric property of the transport coefficient appearing in Eq. (3) can be seen from the Lehmann representation of the correlation function. In the general case we have for the correlation function [1]

$$C^R_{AB}(\nu) = \frac{1}{Z}\sum_{nm}\frac{\langle n|A|m\rangle\langle m|B|n\rangle}{\nu+E_n-E_m+i\eta}\left(e^{-\beta E_n}-e^{-\beta E_m}\right) \tag{S1}$$

where $Z = \mathrm{Tr}[\exp(-\beta H)]$. Therefore, $\mathcal{L}_{AB} \equiv i(\frac{\partial}{\partial\nu}C^R_{AB})\big|_{\nu=0}$ has following form

$$\mathcal{L}_{AB} = -i\frac{1}{Z}\sum_{nm}\frac{\langle n|A|m\rangle\langle m|B|n\rangle}{(E_n-E_m+i\eta)^2}\left(e^{-\beta E_n}-e^{-\beta E_m}\right) \tag{S2}$$

Now if we commute A and B operators, we get

$$\mathcal{L}_{BA} = -i\frac{1}{Z}\sum_{nm}\frac{\langle n|B|m\rangle\langle m|A|n\rangle}{(E_n-E_m+i\eta)^2}\left(e^{-\beta E_n}-e^{-\beta E_m}\right) \tag{S3}$$

Comparing Eq. (S2) and Eq. (S3), gives

$$\Re\mathcal{L}_{AB} = -\Re\mathcal{L}_{BA} \tag{S4}$$
$$\Im\mathcal{L}_{AB} = \Im\mathcal{L}_{BA} \tag{S5}$$

since only real part is giving the current, then one conclude that the transport coefficient is anti-symmetric.

II. DIAGRAMMATIC EXPANSION OF THE CORRELATION FUNCTION AND ITS DERIVATIVE

Fig. S1 shows diagrammatic series expansion of the $C^{(\xi)R}_{\mathbf{j}H'}$. Lines show the full Green's function and $\mathbf{\Gamma}(\mathbf{k}i\omega_m + i\nu_n, \mathbf{p}i\omega_{m'}; \mathbf{p}i\omega_{m'} + i\nu_n, \mathbf{k}i\omega_{m'})$ is irreducible particle-hole interaction vertex. Bare vertexes are given by $\boldsymbol{\lambda}^{(\xi)}_{\mathbf{j}}(\mathbf{k}) = (\partial\mathbf{H}^{(\xi)}_0/\partial\mathbf{k})$ and $\boldsymbol{\lambda}^{(\xi)}_{H'}(\mathbf{k}) = (\partial\mathbf{H}^{(\xi)}_0/\partial\xi)$, respectively. By inserting a $(\partial/\partial i\nu_n)$ vertex and then looking at the $\nu_n \to 0$ limit we can find the perturbation expansion of the $(\partial C^{(\xi)}_{\mathbf{j}H'}/\partial i\nu_n)\big|_{i\nu_n=0}$ (see the second and third row of the Fig. S1). $\boldsymbol{\Lambda}_{\mathbf{j}}$ and $\boldsymbol{\Lambda}_{H'}$ are vertex functions and the last row shows their diagrammatic expansion.

The diagrams contributing at the $(\partial C^{(\xi)}_{\mathbf{j}H'}(\mathbf{q}=\mathbf{0},i\nu_n)/\partial i\nu_n)|_{\nu_n=0}$ can be classified into the following two types: (i) diagrams which are separated into three pieces, each of which has a vertex, by cutting three electron lines, (ii) diagrams which cannot be separated into three pieces by the same operation (See the third row of the Fig. S1).


$C^{(\xi)}_{\mathbf{j}H'}(\mathbf{q}=\mathbf{0},i\nu_n) \equiv$
$(\mathbf{k},i\omega_m+i\nu_n)$
$(\mathbf{p},i\omega_{m'}+i\nu_n)$
$(\mathbf{k},i\omega_m)$
$(\mathbf{p},i\omega_{m'})$
$\lambda_{\mathbf{j}}$
$\lambda_{H'}+$
$\boldsymbol{\Gamma}$
$\cdots$
$\frac{\partial C^{(\xi)}_{\mathbf{j}H'}(\mathbf{q}=\mathbf{0},i\nu_n)}{\partial i\nu_n}\Big|_{\nu_n=0} \equiv$
$= \boldsymbol{\Lambda}_{\mathbf{j}}$
$\boldsymbol{\Lambda}_{H'}+$
$\boldsymbol{\Gamma}'$
$\boldsymbol{\Lambda}_{H'}$
(i)
(ii)
$\boldsymbol{\Lambda}_{\mathbf{j}(H')} =$
$\lambda_{\mathbf{j}(H')}+$
$\boldsymbol{\Lambda}_{\mathbf{j}(H')}$

FIG. S1. (Color online) Diagrammatic expansion of the correlation function, $C^{(\xi)}_{\mathbf{j}H'}(\mathbf{q}=\mathbf{0},i\nu_n)$, and its derivative, $(\partial C^{(\xi)}_{\mathbf{j}H'}(\mathbf{q}=\mathbf{0},i\nu_n)/\partial i\nu_n)|_{\nu_n=0}$. Lines show the full Green's function, $\boldsymbol{\Gamma}$ is irreducible particle-hole interaction vertex and $\boldsymbol{\Gamma}'=(\partial\boldsymbol{\Gamma}/\partial i\nu_n)\big|_{\nu_n=0}$, where ν_n is bosonic frequency. Dash line shows $\partial/\partial i\omega_n$ vertex. $\boldsymbol{\Lambda}_{\mathbf{j}}$ and $\boldsymbol{\Lambda}_{H'}$ are the particle-hole vertex functions and the last row represents their Bethe-Salpeter equation.

III. NON-INTERACTING SYSTEM

Here we show that in the non-interacting case the Eq. (4) reduces to the Berry phase formula. Using the band representation of the Greens function, $\mathbf{g}^{(b)}_{\mathbf{k}\sigma}(i\omega_m,\xi)=[i\omega_m\mathbf{1}-\boldsymbol{\epsilon}^{(\xi)}_{\mathbf{k}\sigma}]^{-1}$, and $\partial_{i\omega_m}\mathbf{g}^{(b)}_{\mathbf{k}\sigma}(i\omega_m,\xi)^{-1}=\mathbf{1}$ one can rewrite Eq. (4) as

$$\begin{aligned}
\frac{d\mathbf{P}^{(\mathrm{el})}}{d\xi} &= -\frac{ie}{2N\beta}\sum_{\mathbf{k}\sigma}\sum_{\omega_m}\varepsilon^{\mathbf{k}\xi}\mathrm{Tr}\Big(\mathbf{g}^{(b)}_{\mathbf{k}\sigma}(i\omega_m,\xi)(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\mathbf{k}})\mathbf{g}^{(b)}_{\mathbf{k}\sigma}(i\omega_m,\xi)(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\xi})\mathbf{g}^{(b)}_{\mathbf{k}\sigma}(i\omega_m,\xi)\Big)\\
&= -\frac{ie}{2N\beta}\sum_{\mathbf{k}\sigma}\sum_{n}\sum_{m}\sum_{\omega_m}\frac{(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\mathbf{k}})_{nm}(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\xi})_{mn}-(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\xi})_{nm}(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\mathbf{k}})_{mn}}{(i\omega_m-\epsilon^{(\xi)}_{n\mathbf{k}\sigma})(i\omega_m-\epsilon^{(\xi)}_{m\mathbf{k}\sigma})(i\omega_m-\epsilon^{(\xi)}_{n\mathbf{k}\sigma})}\\
&= \frac{ie}{2N\beta}\sum_{\mathbf{k}\sigma}\sum_{n}\sum_{m}\frac{\partial}{\partial\epsilon^{(\xi)}_{n\mathbf{k}\sigma}}\sum_{\omega_m}\frac{(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\mathbf{k}})_{nm}(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\xi})_{mn}-(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\xi})_{nm}(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\mathbf{k}})_{mn}}{(i\omega_m-\epsilon^{(\xi)}_{n\mathbf{k}\sigma})(i\omega_m-\epsilon^{(\xi)}_{m\mathbf{k}\sigma})}
\end{aligned}\tag{S6}$$

By summing over Matsubara frequency and using the fact that the Fermi function is one for occupied bands and zero for unoccupied bands, we find (terms with $n=m$ vanish)

$$\frac{d\mathbf{P}^{(\mathrm{el})}}{d\xi}=-\frac{ie}{N}\sum_{\mathbf{k}\sigma}\sum_{n\in occ}\sum_{m\neq n}\frac{(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\mathbf{k}})_{nm}(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\xi})_{mn}-(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\xi})_{nm}(\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\mathbf{k}})_{mn}}{(\epsilon^{(\xi)}_{n\mathbf{k}\sigma}-\epsilon^{(\xi)}_{m\mathbf{k}\sigma})^2}\tag{S7}$$

which reduces to the Berry curvature [2]

$$\begin{aligned}\frac{d\mathbf{P}^{(\mathrm{el})}}{d\xi} &= -\frac{ie}{N}\sum_{\mathbf{k}\sigma}\sum_{n\in occ}\left[\langle\partial_\xi u^{(\xi)}_{n\mathbf{k}\sigma}|\nabla_{\mathbf{k}} u^{(\xi)}_{n\mathbf{k}\sigma}\rangle - \langle\nabla_{\mathbf{k}} u^{(\xi)}_{n\mathbf{k}\sigma}|\partial_\xi u^{(\xi)}_{n\mathbf{k}\sigma}\rangle\right] \\ &= -\frac{ie}{N}\sum_{\mathbf{k}\sigma}\sum_{n\in occ}\left[\partial_\xi\langle u^{(\xi)}_{n\mathbf{k}\sigma}|\nabla_{\mathbf{k}} u^{(\xi)}_{n\mathbf{k}\sigma}\rangle - \nabla_{\mathbf{k}}\langle u^{(\xi)}_{n\mathbf{k}\sigma}|\partial_\xi u^{(\xi)}_{n\mathbf{k}\sigma}\rangle\right]\end{aligned} \tag{S8}$$

using $\langle u^{(\xi)}_{n\mathbf{k}\sigma}|\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\mathbf{k}}|u^{(\xi)}_{m\mathbf{k}\sigma}\rangle = (\epsilon^{(\xi)}_{n\mathbf{k}\sigma} - \epsilon^{(\xi)}_{m\mathbf{k}\sigma})\langle\nabla_{\mathbf{k}}u^{(\xi)}_{n\mathbf{k}\sigma}|u^{(\xi)}_{m\mathbf{k}\sigma}\rangle$ and $\langle u^{(\xi)}_{n\mathbf{k}\sigma}|\frac{\partial\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}}{\partial\xi}|u^{(\xi)}_{m\mathbf{k}\sigma}\rangle = (\epsilon^{(\xi)}_{n\mathbf{k}\sigma} - \epsilon^{(\xi)}_{m\mathbf{k}\sigma})\langle\partial_\xi u^{(\xi)}_{n\mathbf{k}\sigma}|u^{(\xi)}_{m\mathbf{k}\sigma}\rangle$ for $n \neq m$. These expressions can be obtained by taking the $\mathbf{k}(\xi)$-gradient of $\mathbf{H}^{(\xi)}_{\mathbf{k}\sigma}|u^{(\xi)}_{n\mathbf{k}\sigma}\rangle = \epsilon^{(\xi)}_{n\mathbf{k}\sigma}|u^{(\xi)}_{n\mathbf{k}\sigma}\rangle$ and taking the inner product with $\langle u^{(\xi)}_{m\mathbf{k}\sigma}|$. Here, $|\psi^{(\xi)}_{n\mathbf{k}\sigma}\rangle = e^{i\mathbf{k}.\mathbf{r}}|u^{(\xi)}_{n\mathbf{k}\sigma}\rangle$ are the Bloch states and the summation is performed over the occupied bands. The right hand side of Eq. (S8) is the Berry curvature which is gauge invariant and thus observable. [2] The Berry curvature is nonzero in a wide range of materials, in particular, in crystals with broken time-reversal or inversion symmetry. In the periodic gauge [3], the integral on ξ can be done analytically and it leads to a two-point formula for the electric polarization that involves only the initial and final states of the system [4]:

$$\begin{aligned}\Delta\mathbf{P}^{(\mathrm{el})} &= \mathbf{P}^{(\mathrm{el})}(\xi_f) - \mathbf{P}^{(\mathrm{el})}(\xi_i), \\ \mathbf{P}^{(\mathrm{el})}(\xi) &= \frac{ie}{N}\sum_{\mathbf{k}\sigma}\sum_{n\in occ}\langle u^{(\xi)}_{n\mathbf{k}\sigma}|\nabla_{\mathbf{k}}|u^{(\xi)}_{n\mathbf{k}\sigma}\rangle.\end{aligned} \tag{S9}$$

Ionic Hubbard model In 1D-IHM, the energy bands are $\epsilon_\pm(k) = \pm\epsilon(k) = \pm[\Delta^2 + 4t^2\cos^2(ka/2) + 4\xi^2t^2\sin^2(ka/2)]^{1/2}$ and the bare current vertex and the bare distortion vertex in the band representation are given by $(\partial\mathbf{H}^{(\xi)}_0(k)/\partial k) = \mathbf{U}^{-1}(k)[-at\sin(ka/2)\boldsymbol{\tau}_x + a\xi t\cos(ka/2)\boldsymbol{\tau}_y]\mathbf{U}(k)$ and $(\partial\mathbf{H}^{(\xi)}_0(k)/\partial\xi) = \mathbf{U}^{-1}(k)[2t\sin(ka/2)\boldsymbol{\tau}_y]\mathbf{U}(k)$, respectively. $\boldsymbol{\tau}$ are Pauli matrices. $\mathbf{U}(k)$ and $\mathbf{U}^{-1}(k)$ diagonalize the IHM Hamiltonian as $\mathbf{H}_0(k) = \mathbf{U}(k)\boldsymbol{\epsilon}(k)\mathbf{U}^{-1}(k)$ and are given by

$$\mathbf{H}_0(k) = [-2t\cos(ka/2)\boldsymbol{\tau}_x + 2\xi t\sin(ka/2)\boldsymbol{\tau}_y - \Delta\boldsymbol{\tau}_z], \tag{S10}$$

$$\mathbf{U}(k) = \frac{1}{\sqrt{2\epsilon(k)(\epsilon(k)-\Delta)}}[2i\xi t\sin(ka/2)\mathbf{1} + (\epsilon(k)-\Delta)\boldsymbol{\tau}_x + 2t\cos(ka/2)\boldsymbol{\tau}_z], \tag{S11}$$

$$\mathbf{U}^{-1}(k) = \frac{1}{\sqrt{2\epsilon(k)(\epsilon(k)-\Delta)}}[-2i\xi t\sin(ka/2)\mathbf{1} + (\epsilon(k)-\Delta)\boldsymbol{\tau}_x + 2t\cos(ka/2)\boldsymbol{\tau}_z]. \tag{S12}$$

At small ξ the denominator of Eq. (S7) is given by $(\epsilon^{(\xi)}_{-k\sigma} - \epsilon^{(\xi)}_{+k\sigma})^2 \simeq 4(\Delta^2 + 4t^2\cos^2(ka/2))$ while the imaginary part of the numerator is $(\partial\mathbf{H}^{(\xi)}_0/\partial k)_{12}(\partial\mathbf{H}^{(\xi)}_0/\partial\xi)_{21} - (\partial\mathbf{H}^{(\xi)}_0/\partial\xi)_{12}(\partial\mathbf{H}^{(\xi)}_0/\partial k)_{21} \simeq 2t^2\sin^2(ka/2)\cdot(\Delta/\sqrt{\Delta^2 + 4t^2\cos(ka/2)})$. Therefore, the polarization at small ξ is given by

$$P^{(\mathrm{el})}_{\Delta,U}(\xi) \simeq -\frac{2ea}{N}\sum_k\frac{(\Delta/t)\sin^2(ka/2)}{[(\Delta/t)^2 + 4\cos^2(ka/2)]^{3/2}}\xi. \tag{S13}$$

---

[1] H. Bruus and K. Flensberg, *Many-Body Quantum Theory in Condensed Matter Physics: An Introduction* , Oxford University Press (2004).

[2] D. Xiao, M. C. Chang, and Q. Niu, Rev. Mod. Phys., **82**, 1959 (2010).

[3] In periodic gauge the Brillouin zone has the topology of a torus: the two points $\mathbf{k}$ and $\mathbf{k}+\mathbf{K}$ can be identified as the same point, where $\mathbf{K}$ is the reciprocal lattice vector. From boundary condition $|\psi^{(\xi)}_{n\mathbf{k}\sigma}\rangle$ and $|\psi^{(\xi)}_{n\mathbf{k}+\mathbf{K}\sigma}\rangle$ can at most differ by a phase factor. The torus topology is realized by making the phase choice $|\psi^{(\xi)}_{n\mathbf{k}\sigma}\rangle = |\psi^{(\xi)}_{n\mathbf{k}+\mathbf{K}\sigma}\rangle$. Consequently, $|u^{(\xi)}_{n\mathbf{k}\sigma}\rangle$ and $|u^{(\xi)}_{n\mathbf{k}+\mathbf{K}\sigma}\rangle$ satisfy the following phase relation; $u^{(\xi)}_{n\mathbf{k}\sigma}(\mathbf{r}) = \exp(i\mathbf{K}.\mathbf{r})u^{(\xi)}_{n\mathbf{k}+\mathbf{K}\sigma}(\mathbf{r})$.

[4] R. D. King-Smith and D. Vanderbilt, Phys. Rev. B **47**, 1651 (1993).